



A service of the National Library of Medicine and the National Institutes of Health

My NCBI [?] [Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0

1: Am J Epidemiol. 1981 Aug;114(2):209-17.

Full Text Links
OXFORD JOURNALS

Breast cancer incidence in women with a history of progesterone deficiency.

Cowan LD, Gordis L, Tonascia JA, Jones GS.

In order to investigate the nature of the association of involuntarily delayed first birth and risk of breast cancer, 1083 white women who had been evaluated and treated for infertility from 1945-1965 were followed prospectively through April 1978 to ascertain their breast cancer incidence. These women were categorized as to the cause of infertility into two groups, those with endogenous progesterone deficiency (PD) and those with nonhormonal causes (NH). Women in the PD group had 5.4 times the risk of premenopausal breast cancer compared to women in the NH group. This excess risk could not be explained by differences between the two groups in ages at menarche or menopause, history of oral contraceptive use, history of benign breast disease or age at first birth. Women in the PD group also experienced a 10-fold increase in deaths from all malignant neoplasms compared to the NH group. The incidence of postmenopausal breast cancer did not differ significantly between the two groups.

PIP: In order to investigate the nature of the association of involuntarily delayed 1st birth and breast cancer risk, 1083 white women who had been evaluated and treated for in fertility from 1945-65 were followed prospectively through April 1978 to ascertain their breast cancer incidence. These women were categorized as to the cause of infertility into 2 groups, those with endogenous progesterone deficiency (PD) and those with nonhormonal causes (NH). Women in the PD group had 5.4 times the risk of premenopausal breast cancer as compared to women in the NH group. This excess risk could not be explained by differences between the 2 groups in age at menarche or age at menopause, history of oral contraceptive use, history of benign breast diseases, or age at 1st birth. Women in the PD group also experienced a 10-fold increase in deaths from all malignant neoplasm compared to the NH group. The incidence of postmenopausal breast cancer did not differ significantly between the 2 groups. author's modified

PMID: 7304556 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort By Send to

Related Links

Breast cancer among young U.S. women in relation to oral contraceptive use. [Am J Epidemiol. 1994]

Risk factors for breast cancer according to family history of breast cancer. For the Nurses' Health Study. [Breast Cancer. 1996]

Age-specific patterns of association between breast cancer and risk factors in black women, ages 20 to 39 and 40 to 54. [Ann Epidemiol. 1994]

Cancer incidence in a cohort of infertile women. [Am J Epidemiol. 1987]

Breast cancer in premenopausal and postmenopausal women. [Breast Cancer Inst. 1974]

See all Related Articles...

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Privacy Statement | Freedom of Information Act | Disclaimer